

Modeling physical and chemical pollution of Sebou river waters

Kenitra, Morocco

SIBARI MOHAMED, HAMDAOUI FARID, LAKHLIFI MOSTAPHA, ELATMANI AYOUB,
ACHHAR ABDERRAHMANE, KHADIJA EL KHARRIM, DRISS BELGHYTI

Department of Biology
University Ibn Tofail
PB: 133, Campus Universitaire, 14000 Kénitra
MOROCCO

Abstract: - The physicochemical characterization of the raw surface waters of the lower Sebou river revealed that this river is heavily loaded with mineral matter. Thus, for the electrical conductivity (EC) a wide variation in the chemical composition of water, it varies between a minimum of 629 $\mu\text{S} / \text{cm}$ and a maximum of 22760 $\mu\text{S}/\text{cm}$. The average pH is between 8.00 and 8.77. The pH is basic but remains acceptable according to irrigation standards. Nitrates range from 0.04 to 2692 mg / L. Chlorides range from 145.55 to 860.27 mg / L. Sulphates range from 37.62 to 441.4 mg / L. Sodium ranges from 51 to 2530 mg / L.

It can be deduced that the lower Sebou sub-basin, whose mineralization faithfully follows the levels of dissolved salts, salinity, chlorides, sodium and potassium, is subject to the different types of pollution of natural origin, which are mainly mineral by (dissolving natural karst limestone substrates, Atlantic tidal, ocean spray) and anthropogenic (agricultural, industrial and urban).

In conclusion, it appears from this study that the waters of the lower Sebou waters are too polluted and we recommend a proper treatment of domestic and industrial wastewater to reduce the nuisance that this river undergoes and also to alleviate the loss of this much-coveted and sought-after water resource..

Key-Words: - Sebou Waters; Hydraulics, Physico-chemistry, Pollution, Kenitra, Morocco.

1 Introduction

Water quality is defined by physical, chemical and biological parameters, but also by its use. Thus, water unfit for human consumption can be adapted to irrigation, fish farming or to cool industrial circuits [1-3]. The rational management of water resources in the Kenitra Gharb area has become the main issue for local decision-makers to adopt a fair policy and which takes into consideration the importance of this resource and the challenge of increasing water resources. The Sebou river and its tributaries drain an area of 34000 km². It extends for more than 600 km starting in the Middle Atlas under the name of Guigou river. It opens in the Atlantic to Mehdia, through its estuary 35 km in length. The rise of marine waters being stopped at the level of the guard dam, immediately downstream of Sidi Allal Tazi city [4]. In addition, the Sebou river is home to many pollutant spills from a variety of sources. The Sebou watershed, an extremely important area from a socio-economic point of view, is one of the most affected areas in Morocco. The existence of two of the main agricultural plains of the country as well as the multitude and diversity of

industrial units and urban wastewater effluents in the major cities of the basin (Fez, Allal Tazi, Mechraa Bel Ksiri, Dar Gueddari, Kenitra), not to mention the uncontrolled dumping of household waste, which are the main causes of the deterioration of the quality of Sebou waters.

In our present study it is proposed to examine the physicochemical surface water of the lower Sebou sub-basin. This characterization of the levels and concentrations of the organic and mineral loads of Sebou raw water consists of a monitoring of the pH, EC electrical conductivity, sodium, chloride, sulphate, calcium, magnesium, potassium, bicarbonate, ammonium and nitrates.

2 Material and Methods

2.1 Study area

The Gharb region is bordered on the west by the Atlantic sea, bordered to the north by the pre-Rif hills and to the south by the Maâmora shelf (**Fig.1**). It is composed of a coastal zone (dune cord, flooded

depressions, interior dunes), continental borders and the central alluvial plain of the lower Sebou which is the main river. The Sebou basin forms a basin between the Rif in the North, the Middle Atlas and the Meseta in the South, the Taza corridor in the East and the Atlantic sea in the West. It is the most important basin of the kingdom with approximately 38380 km² and currently contains a total population of 5.73 million inhabitants, of which 49% in urban and 51% in rural areas. It is characterized by an agricultural and industrial activity that contributes significantly to national economy.

The climate prevailing on the whole basin is of Mediterranean type with oceanic influence and inside the basin the climate becomes more continental. The Sebou basin has a very developed industrial activity. Large units at the basin scale are: sugar mills, paper mills, oil mills, tanneries, cement plants, the textile industry and the oil refinery. The taking of a water sample is a delicate operation to which the greatest care must be taken, it determines the analytical results and the interpretation that will be given. In general, the sample must be homogeneous and representative, and not modify the physicochemical characteristics of the water (dissolved gas, suspended matter, etc) [5]. Sampling equipment should be given special attention. The washing of the flasks will depend on the desired analyzes on the sample. The most frequently used sampling method is instant sampling. The vials are filled without shaking the water and sometimes without contact with the air [6-7].

2.2. Water analysis

1000 ml polyethylene bottles were previously rinsed with distilled water and then with the sample water in the field. Sampling was done in areas where the water is not stagnant and in the direction of flow. It is carried out in total immersion, so that the bottles are filled flush without air bubbles, in order to minimize the contamination on the one hand, and the evolution of the samples on the other. The water samples taken for analysis were transported at low temperature (4 °C) in portable coolers to the laboratory where analyzes were carried out. In addition, from one campaign to another, the samples were taken at approximately the same time and place for the same station.

In the present study the parameters that were analyzed are: pH, electrical conductivity (EC), calcium (Ca⁺⁺), magnesium (Mg⁺⁺), sodium (Na⁺), potassium (K⁺), carbonate and bicarbonate (CO₃⁻⁻, HCO₃⁻), chlorides (Cl⁻), sulphates (SO₄⁻⁻), ammonium (NH₄⁺) and nitrate (NO₃⁻). The devices

used are Assays C831, Jenway flame photometer, NOVASPEC II pharmacy-type spectrophotometer, UV-Visible spectrophotometer [8-11].

Calcium and magnesium are determined by complexometry with EDTA in the presence of Eriochrome black T. Determination of carbonates and bicarbonates by a solution of 0.02N sulfuric acid in the presence of phenolphthalein and bromocresol green as a colored indicator. Determination of the combined chloride in the chloride state by silver nitrate, in the presence of a solution of potassium chromate. Determination of sulfates by colorimetry by precipitation of sulphate ions in the presence of barium chloride in a hydrochloric acid medium in the form of barium sulphate. Determination of nitrates and ammoniums by distillation in the presence of a catalyst respectively magnesium oxide and alloy DEVARDA. NH₄⁺ and NO₃⁻ are collected in a boric acid solution and finally assay with H₂SO₄.




Fig. 1: Localization of sampling Sebou basin.

3 Results and Discussion

The evaluation of raw water pollution of the lower Sebou was made according to the determination of a certain number of physicochemical parameters characterizing the waters. In the light of this work which contributes to enriching the bases of the data accumulated on the Sebou basin, and to make it possible to clarify the degree of its pollution thanks to the results obtained during the period of our internship within the Regional Office of implementation agricultural value of Kenitra.

It can be deduced from **Tables 1, 2 and Figure 2**, that the sub basin of the lower Sebou river is subject to different types of pollution of natural origin which are mainly mineral (by dissolution of the natural substrate, Atlantic tides) and anthropogenic (agricultural, industrial and urban).

The thermal regime of the Sebou hydrographic network follows that of the Mediterranean climate cold in November and warmer in summer.

The pH does not show any significant variation and the waters are generally alkaline ranging between 8.0 and 8.77 (**Tab.1**) following the crossing of limestone and marl-limestone soil characterizing the basin.

Mineralization accurately follows dissolved salt salinity, chloride, sodium and potassium level (**Tab. 1, 2 ; Fig.2**). It results essentially from the leaching of the karstic limestone and kelp-like terrain and ocean spray. Indeed, the electrical conductivity that reflects salinity (Tab.2) varies from 629 to 22760 $\mu\text{S} / \text{cm}$ and far exceeds the Moroccan irrigation standard ($> 2700 \mu\text{S} / \text{cm}$) [12-14].




Fig. 2: Piper Diagram of water of Sebou river

Table 1: Physicochemical of the raw waters of Sebou river.

Stations	pH	NO ₃ - mg/L	CL - mg/L	SO ₄ - mg/L	HCO ₃ - mg/L	CO ₃ - mg/L
S1	8,62	9,3	213	314,64	233,02	12
S2	8,65	10,42	161,88	159,18	214,72	18
S3	8,39	0,24	202,35	150,25	213,5	6
S4	8,46	17,11	154,78	141,18	275,72	0
S5	8	63,36	243,53	181,94	246,44	0
S6	8,12	20,58	202,35	151,62	362,34	0
S7	8,69	86,92	248,5	183,59	241,56	21,6
S8	8,33	188,6	385,53	258,32	323,3	13,2
S9	8,49	827,9	230,4	106,7	75,64	0
S10	8,24	2692	860,27	113,1	122	0
S11	8,4	260,8	269,09	37,62	100,04	0
S12	8,31	886,9	476,41	276,57	84,18	0
S13	8,77	94,6	461,31	359,29	246,44	49,2
S14	8,21	162,2	397,7	441,4	178,12	42
S15	8,73	59,9	304,59	248,5	241,56	18
S16	8,33	693,8	145,55	54,04	108,58	0

Concerning the nitrate contents (**Tab.1**), the values oscillate between 0.24 mg / L and 2692 mg / L and clearly translate the pollution of agricultural origin by the nitrogenous fertilizers, the wastewater and leachates of the wild discharges [15-21].

The Piper diagram (**Fig.2**) shows that globally the waters of the lower Sebou are hyper-chlorinated calcium, hyper-sulphated calcium, chlorinated sulphated calcium and magnesium or chlorinated sodium and potassium or sulphated sodium [22, 23].

Moreover, the projection of physicochemical data in the Wilcox diagram and Wilcox Log diagram, shows that the quality of the waters of the lower Sebou varies between Poor and Bad and rarely excellent and especially have a degraded quality because the alkalinizing power of sodium (SAR). The waters of the lower Sebou are classified in the group C3S4 and C4S4 and are unsuitable for irrigation [24-25].

Table 2: Physicochemical of the raw waters of Sebou river

Stations	Ca ²⁺ mg/L	Mg ²⁺ mg/L	K ⁺ mg/L	Na ⁺ mg/L	NH ₄ ⁺ mg/L	TH mg/L	CE	References:
S1	97,6	89,04	6,44	1240	0,18	6,15	1240	[1] C. Boutin and N. Dias, “Impact de l'épandage des eaux usées de la ville de Marrakech sur la nappe phréatique”, Bull. Fac. Sci. Marrakech (Sect. Sci. Vie), 3 (1987), pp. 5-25.
S2	118	46,56	6,24	1270	0,43	4,89	1240	
S3	126,4	59,52	2,54	1560	0,04	5,72	1240	
S4	166,4	17,76	5,27	1360	0,68	4,89	1240	[2] M. Benyakhlef, S. Naji, D. Belghyti, Y. El guamri et T. Hassouni, “ Qualité de l'eau de boisson dans la région du Gharb (Kénitra, Maroc)” 2011.
S5	150,8	44,4	4,29	1820	0,22	5,82	1240	
S6	169,2	68,64	9,56	1330	0,18	7,09	1400	
S7	148,4	51,36	4,68	1470	0,68	5,85	1400	[3] H. Abouzid et A. Outair, “ Les Nitrates dans les eaux”, 7ème Congrès Mondial des ressources en eau, Rabat, Maroc, 13-18 Mai 1991, Volume 2.
S8	220,4	111,6	12,48	1840	0,5	10,16	1400	
S9	217,6	17,28	3,71	51	0,68	6,16	629	[4] ORMVAG, “Office Régional de mise en Valeur Agricole Charb Maroc : Etude pédologique au 1/20 000 de la Troisième Tranche d'Irrigation (TTI) sur une superficie de 100.000 ha. Zone M'nasra, Z1-Z2. Kénitra, Maroc”, Rapport inédit, 1994, 180p.
S10	914,4	631,2	5,07	140	1,76	49,82	629	
S11	148,4	26,64	6,63	190	15,34	4,82	629	
S12	314,8	39,36	17,55	390	0,54	9,82	629	
S13	170,8	124,08	9,75	2530	2,66	9,44	2200	[5] J. Rodier, “L'analyse de l'eau: eaux naturelles, eaux résiduaires, eau de mer : physico-chimie, bactériologie et biologie”, Ed. Dunod, Paris, France, 8 (1996), 1383p.
S14	1072,8	74,88	15,99	150	1,26	57,82	2200	
S15	144,8	94,08	7,41	400	1,29	7,34	1664	
S16	174	99,36	2,73	120	1,51	8,49	880	

4 Conclusion

Adjacent agricultural activities occur well in the waters of the Lower Sebou sub-basin by significant concentrations of nitrates and sulphates which enter the water stream by runoff and leaching of nitrogenous and phosphorus fertilizer and phytosanitary products [26-27]. The upstream-downstream distribution of physicochemical parameters, reflects deteriorated situations of water quality in salts and chlorides in relation to the rise of marine saline waters.

The present work has revealed the poor quality of the waters of the lower Sebou but remains incomplete and needs to be deepened by analyzes of trace heavy metals and pesticides to provide the scientific and technical bases for decision-makers [28-30].

- [7] O.N.E.P, “Méthodologie d'analyse de l'eau au laboratoire. Mode opératoire: Contrôle de la pollution des eaux destinées à l'alimentation en eau potable”, 1998.
- [8] L. Matini, J.M. Moutou et M.S. Kongo-Mantono, “ Evaluation hydro-chimique des eaux souterraines en milieu urbain au Sud-Ouest de Brazzaville, Congo”, 2012.
- [9] T. El Hammoumi et Belghyti D., “Caractérisation physicochimique des eaux potables Produit Par la station de traitement de Mkansa” (Maroc, 2012).

- [10] L. Bentouati et Bouzidi A., "étude de la qualité des eaux souterraines de la wilaya de Sétif, algésaire", Journal Scienceslip, 2011.
- [11] B. Benkabbour, "Exploration, évaluation et protection des ressources hydriques en zones côtières Marocaines : Approche Géophysique, Hydro chimiques, modélisation et S.I.G: Cas de la Maàmora occidentale (Bassin du Rharb-Maàmora)", Thèse de Doctoral National, Université Ibn Tofail, 2002.
- [12] Administration Hydraulique (AH), "Etat de la qualité des ressources en eaux dans le bassin du Sebou, année 1989-1990", Ministère des Travaux Public, de la Formation Professionnelle et de la Formation des Cadres. 1996.
- [13] S. Akhiar, "Caractérisation des eaux souterraines de la ville de Mechraa Bel Ksiri". Mémoire Master Eaux usées. Université Ibn Tofail, Kénitra, 2009.
- [14] M. Hilali, "Hydrogéologie et modélisation de l'intrusion marine dans les aquifères côtiers de Martile et de Sahel- Maroc", Thèse de Doctorat en Sciences Appliquées. Université. Mohammed V-Agdal, Ecole Mohammedia d'ingénieurs, 2002, 158p.
- [15] Secrétariat d'Etat chargé de l'Environnement, "Etude pour programme d'action visant à minimiser et à contrôler l'impact des engrais et des pesticides sur l'environnement du bassin de Sebou", (Projet de l'environnement du Sebou), Secrétariat d'état chargé de l'environnement, (Maroc), (1999), 43p.
- [16] Z. Saadi, A. Maaslouhi, M. Zeraouli et J. P Gaudet, "Analyse et modélisation des variations saisonnières des concentrations en nitrates dans les eaux souterraines de la nappe Mnasra, Maroc", C. R. Acad. Sci., Sér. 2, Sci. Terre Planètes, 329, 8, 1999, pp. 579-586.
- [17] MPCI, "Impact environnemental de l'usage des eaux usées d'assainissement dans l'irrigation des agricultures", ministère de la planification et de la coopération internationale, juillet 2005.
- [18] A. Alemad , Nagi M., Ibeda A., Nasser R., Alwathaf Y., Elrhaouat O., Elkharrim K., Babaqi A., Belghyti D., "The impact of sana'a solid waste on the quality of groundwater in Yemen", 2nd International Conference on Water and Society, 4 - 6 September 2013, New Forest, UK, Paper DOI: 10,2495/WS130151.
- [19] M. S. Coyne and J. M. Howell, "Agricultural Impacts on Fecal Contamination of Shallow Groundwaters in the Bluegrass Region of Kentucky", Soil Science News and Views, 15, 6, 1994, pp. 1-3.
- [20] A., B. Krira. Chakour et H. Fouta, "Intensification de l'agriculture et son impact sur l'environnement. Cas des nitrates dans la nappe phréatique de M'nasra du Gha", Actes 1er Colloq. Sur le Développement agric. Rech. Agron. Au niveau de la région du Ghab, 2001.
- [21] M., Zeraouli, "Pollution par les nitrates. Premiers résultats de la situation actuelle dans la nappe des Mnasra" (décembre 1992-janvier 1993) », Office régional de mise en valeur agricole du Gharb, Département de développement agricole, Service des études de développement agricole, Bureau Agro- Pédologique, Publication interne ORMVAG, septembre, 1993.
- [22] L. Zilliox, C.Schenc, H. Kobus et B. Huwe, "Pollution par les nitrates: Quels remèdes ? Supplément", La Recherche Suppl. les enjeux de l'agriculture en Europe, 227, 1990, pp. 18-21.
- [23] K.. El Bouqdaoui, Aachib, M., Blaghen, M., et Kholtei, S., "Modélisation de la pollution par les nitrates de la nappe de Berrechid, au Maroc", 2009.
- [24] L. Zouhri, "Structure et modélisation hydrodynamique de l'aquifère de la Maamora (Maroc)". Thèse, Univ. LilleI, 2000, 218p.
- [25] M. Nisbet et Verneaux J., "Composantes chimiques des eaux courantes : discussion et proposition de classes en tant que bases d'interprétation des analyses chimiques", Annls Limnol, 6, 2, 1970.
- [26] O.N.E.P, "Alimentation en eau potable, Menaces de pollution", 1999.
- [27] Laferriere. M, J. J. Minville, J. Lavoie et P. Payment, "L'industrie porcine et les risques reliés à la santé humaine", Bull. Information Santé Environnem, Québec, 7, 2, 1996, pp. 1-4.
- [28] Organisation Mondiale de la Santé (O.M.S.), "Charte d'Ottawa pour la promotion de la sant", Copenhagen, Bureau régional de l'Europe, (1986).

[29]. CSE. Conseil Supérieur de l'Eau, “Aménagement optimal des eaux de l'oued Ouergha: Réalisation du barrage Mjara”, Rabat, Maroc. 1988.

[30] A.B.H.S., “Etude d'actualisation du plan directeur d'aménagement integer des resources en eau du bassin hydraulique du Sebou. Note de synthèse”, Septembre 2011, Agence du Bassin Hydraulique du Sebou, Fès, Maroc.