



























- [43] Jaramillo J. and Vega F.E., *Aphanogmus* sp. (Hymenoptera: Ceraphronidae): a hyperparasitoid of the coffee berry borer parasitoid *Prorops nasuta* (Hymenoptera: Bethyridae) in Kenya, *Biocontrol Science and Technology*, 19, 2009, pp. 113-116.
- [44] Ticheler J.H.G., Etude analytique de l'épidémiologie du scolyte des grains de café, *Stephanoderes hampei* Ferr., en Côte d'Ivoire, Mededelingen Landbhoogeschool Wageningen, Netherlands, 61, 1961, pp. 1-49.
- [45] Borbón-Martínez M.O., Bio-écologie d'un ravageur des baies de caféier *Hypothenemus hampei* Ferr. (Coleoptera: Scolytidae) et de ses parasitoïdes au Togo, Thèse de Doctorat Ph.D, Université Paul Sabatier, Toulouse, France, 1989, pp. 1-185.
- [46] Bustillo A.E., Bernal M.G., Benavides P. and Chavez B., Dynamics of *Beauveria bassiana* and *Metarhizium anisopliae* infecting *Hypothenemus hampei* (Coleoptera: Scolytidae) populations emerging from fallen coffee berries, *Florida Entomology*, 82, 1999, 491-498.
- [47] Graziani L., Déterminisme des sex-ratios primaires et secondaires chez les vertébrés supérieurs : Rôle du milieu physique et social des individus. Diplôme d'Etude Approfondie Analyse et Modélisation des Systèmes Biologiques, Rapport bibliographique, UCB Lyon1, 1994, pp. 1-38.
- [48] Paul A. and Thommen D., Timing of birth, female reproductive success, and infant sex ratio in semifree-ranging Barbary macaques (*Macaca sylvanus*). *Folia Primatology*, 42, 1984, pp. 2-16.
- [49] Voula V.A., Manga E.F., Messi A.L., Mahob J.R. and Begoude B.A., Impact of mirids and fungal infestation on dieback of cocoa in Cameroon, *Journal of Entomology and Zoology Studies*, 6(5), 2018, pp. 240-245.
- [50] Roeder K.D., An experimental analysis of the sexual behavior of the praying mantis (*Mantis religiosa* L.), *The Biological Bulletin*, 69(2), 1935, pp. 203-220.
- [51] Hurd L.E., Eisenberg R.M., Fagan W.F., Tilmon K.J., Snyder W.E., Vandersall K.S., Datz S.G. and Welch J.D., Cannibalism reverses male-biased sex ratio in adult mantids: female strategy against food limitation. *Oikos*, 1994, pp. 193-198.
- [52] Mbondji M., Knowledge to date on *Stephanoderes hampei* bionomics Coleoptera Scolytidae, *Annale Faculté des Sciences, Université Yaoundé I, Cameroun*, 17, 1974, pp. 95-104.
- [53] Vega F.E., The rise of coffee, *American Scientist*, 96, 2008, pp. 138-145.
- [54] Wintgen J.N., *Coffee: Growing, processing, sustainable production*, Wiley-VCH Edition, 2004.
- [55] Bosselmann A.S., Dons K., Oberthur T., Olsen C.S., Ræbild A. and Usma H., The influence of shade trees on coffee quality in small holder coffee agroforestry systems in Southern Colombia. *Agriculture, Ecosystems and Environment*, 129, 2009, pp. 253-260.
- [56] Jaramillo J., Torto B., Mwenda D., Troeger A., Borgemeister C., Poehling H.M. and Francke W., Coffee berry borer joins bark beetles in coffee klatch, *PLoS One*, 8, 2013, pp. e74277.
- [57] Mariño Y.A., Pérez M.E., Gallardo F., Trifilio M., Cruz M. and Bayman P., Sun vs. shade affects infestation, total population and sex ratio of the coffee berry borer (*Hypothenemus hampei*) in Puerto Rico, *Agriculture, Ecosystems and Environment*, 222, 2016, pp. 258-266.
- [58] Salazar H.M., Aristizábal L.F. and Mejía C.G., Investigación participativa en relación con el manejo de la broca del café *Hypothenemus hampei* (Coleoptera: Scolytidae) en el proceso de beneficio. *Revista Colombiana de Entomología*, 29, 2003, pp. 57-62.