Efficacy of biogenic synthesised silver nanoparticles on behavioural changes of *Helicoverpa armigera*

ISHWARYA M S¹, LOKESH S²* ^{1,2}Department of Biotechnology, University of Mysore, Mysuru-570 005, Karnataka, INDIA

Abstract: The grounds of this study were to utilize the biogenic synthesized silver nanoparticles from *Leucas aspera, Cyperus rotundus and Cynodon dactylon* for the control of polyphagous pest *Helicoverpa armigera*. Synthesized silver nanoparticles were subjected for different characterization like XRD, EDX, FTIR and Raman spectroscopy. The bioassays (i.e. larval weight and survival rate) and also its interaction with gut protease activity (trypsin and Chymotrypsin) was evaluated employing LA, CR, and CD AgNPs, as well as crude plant extracts against the 4th instar larvae. The stretching vibration of distinct functional groups of the samples was demonstrated by the existence of FTIR measurements. XRD graphs confirm the AgNPs crystallize in centered cubic phase and the change in peak intensity with different plant extracts was noticed. The body mass, length of the larvae were dramatically lowered by the synthesized sample of silver nanoparticles. They also suppressed the activity of gut proteases (trypsin and chymotrypsin). Our studies could aid in the development of new approaches to address insecticidal resistance in *H. armigera*.

Key words: *Helicoverpa armigera*, Biosynthesized silver nanoparticles, *Leucas aspera*, *Cynodon dactylon*, *Cyperus rotundus*, Antifeedant, Larvicidal, gut protease.

Received: August 15, 2022. Revised: October 26, 2022. Accepted: November 28, 2022. Published: December 29, 2022.

1. Introduction

Agriculture is by far the most basic aspect of every country and also an integral aspect in progressing toward an immobile human lifestyle. To compensate for crop losses and financial constraints caused by numerous biological and chemical variables, the farming community have undergone substantial changes. Chemical insecticides and fertilisers were used extensively to boost vields. Regrettably, unrestrained use of chemical insecticides and fertilisers has caused a spike in pathogen and pest resistance, reduced soil species diversity, reduction in nitrogen fixation contributes to pesticide bioaccumulation, pollinator decline, destruction of bird habitat, toxic and nonbiodegradable, negatively affecting human health [Oliveira et al., 2014; Lade et al., 2017; Randall et al., 2013]. Despite the fact that assessing the chemistry of bio pesticides offers a reasonable framework based on evolutionary selection for chemicals that are more specific to an insect, have faster decomposition, lower environmental persistence and don't often increase in concentration in the environment. Biopesticides have been the least preferred by farmers because their efficacy is poor at different geographical conditions and laggard in pest control [Kamaraj et al., 2008; Kamaraj et al., 2008; Senthil, 2015].

Nanoparticles could be an excellent and viable choice for achieving long-term agricultural success. The dexterity of nanotechnology has the potential to precisely form matter of 10-100nm size i.e. small size can increase the surface area of active compound intrun increase solubility, improving bioavailability (i.e. increased contact with the target cell) [Sasson et al., 2007; Campolo et al., 2017; Gruère et al., 2011], safeguard the active component from degradation so that it can be released for a long and with precision, efficacy. time and specificity [Worrall et al., 2018]. Plant-based silver nanoparticles production is a one-step biosynthesis approach that is speedy, low-cost and eco-friendly [Huang et al., 2011]. The core objective of the green synthesized silver nanoparticles strategy is to provide larvicidal, insecticidal. antibacterial. viral and inflammatory characteristics. Various experts from all over the world have cited multiple studies demonstrating the insecticidal activity of AgNPs. The physiological effect of AgNPs derived from Cassia fistula L aqueous extracts was investigated using A. albopictus and Culex pipiens fourth instar larvae and also enzyme assays supported the study [Fouad et al., 2018].

Cotton, pigeon pea, chickpea, tomato, okra, black gram, corn, and a range of other crops are all affected by *Helicoverpa armigera*. The pest's larvae feed on the reproductive systems of plants, causing a loss of around a one third of the yield. H. armigera has the ability to adapt to a variety of cropping systems due its high polyphagous fecundity, and pesticide resistance. proteases such trypsin Serine as and chymotrypsin are prevalent in the digestive tract of H. armigera [Srinivasan et al., 2006]. Insect gut proteases are crucial for the energy and growth of insects by aiding in the digestion of proteinaceous foods. One of the most important techniques for reducing the insect pest is to For inhibit gut proteases. sustenance, Lepidopteran insect pests depend mainly on trypsin and chymotrypsin-like digestive proteases [Sharma et al., 2000; Swathi et al., 2016]. The identification of gut protease inhibitors from plants that are effective against H. armigera has indeed been described by several experts. Using biosynthesized gold nanoparticles from Jatropha curcas L. latex, serine protease restriction can be seen in a range of insects, including Aedes aegypti, beetles and Pseudococcidae pests. [Patil et al., 2016]. AgNPs were observed to kill juvenile Hematophagous flies, Hippobosca maculate and bovine ticks. Rhipicephalus (Boophilus) microplus, across both high and low quantities. [Santhoshkumar 2012]. et al., Culex quinquefasciatus and Anopheles subpictus 4th instar larvae were successfully controlled using AgNPs synthesised from *Eclipta prostrata* leaf extract [Rajakumar et al., 2011].

The present study aims to utilise Leucas aspera, Cyperus rotundus, and Cynodon dactylon which are exquisite plant species (i.e. most ecological, locally available, reproducible, economically priced). They compete with most crops for water, sunlight, nutrients, and space, making it an excellent plant for fabrication of silver nanoparticles. XRD, EDX, FTIR and Raman spectroscopy were used to characterise the synthesised silver nanoparticles. The behavioural (i.e. larval weight and survival rate) alterations in H. armigera were detected with AgNPs and leaf extract and also evaluating the alter efficacy of gut protease activity in H. armigera (trypsin and Chymotrypsin).

2. Materials and Procedures

The silver nitrate used in this study was purchased from Sigma-Aldrich. Plant species such as *Leucas aspera*, *Cyperus rotundus*, and *Cynodon dactylon* were randomly picked from the University of Mysore in Mysuru, India.

3. Preparation of plant extracts and biogenic synthesis of silver nanoparticles

All three plants were cleansed by washing them in the double distilled water. Thereafter, the material was dried in the shade. In a magnetic stirrer, the powdered leaves were blended with 100 mL distilled water and incubated/heated for 30 minutes. The extract was refrigerated and decanted before being filtered and kept at -4°C using Whatman filter paper. LA, CR, and CD were assigned to crude leaf extracts *from L. aspera, C. rotundus, and C. dactylon*, respectively.

Mixing 50 mL of plant extract in 100 mL of 1 mM AgNO₃ solution at 70°C while stirring constantly yield silver nanoparticles. The colour variations in the reaction mixture from pale yellow to reddish brown indicate the production of AgNPs. Followed by next step, AgNPs formed were washed 5-6 times with distilled water and ethanol before being centrifuged for 10 minutes at 10000 RPM, pellet obtained was freeze, dried and stored at -4°C until use. The synthesised AgNPs were subjected to UV-vis, for characterization. AgNPs produced with *L. aspera, C. rotundus, and C. dactylon* plant extract were designated as LA-AgNPs, CR-AgNPs, and CD-AgNPs, respectively.

3.1. Characterization Techniques

The morphological and structural analysis of synthesised AgNPs was performed using Fourier Transform infrared spectroscopy, X-ray Dispersive diffraction. Energy X-Rav Spectroscopy and Raman spectroscopy. An Xray diffractometer was used to determine the characterisation structural and crystalline structure of metal nanoparticles (Rigaku, BD63000074). FTIR spectrophotometer is used to examine the chemical compositions of silver the nanoparticles and also existence of (Nicolet-6900, functional groups Model: 912A0637). HORIBA's Xplora Plus-42308 was used to analyse the Raman spectra of silver nanoparticles.

3.2 Tested insects

The culture of the tested insect *H. armigera* (4th instar larvae) was received from an insect breeding lab, ICAR-NBAIR, Hebbal Bengaluru 560024. They were raised in a lab at a temperature of 26°C, a relative humidity of 65.5°RH and 12 hours of photoperiod on a

chickpea-based artificial diet. Chickpea (100g), methyl PHB (2g), sorbic acid (1g), yeast (10g), agar (12.75g), ascorbic acid (3.25g), multivitamin tab (1 capsule), vitaminE (2 capsule), streptomycin (0.25g), 10% formalin (5ml), and water made up the artificial diet. The larvae of the 4thinstar were collected and tested in the lab in a bioassay.

3.3.Bioassay of *Helicoverpa armigera* using silver nanoparticles

Antifeedant assay and Larvicidal activity: diet choice tested with required The modification was used to determine the antifeedant larvicidal activity and in Helicoverpa armigera 4th instar larvae. In brief, the larvae were divided into 3 groups of 4 larvae in each group and the selection was complete randomized design. All weights were measured using a monopam balance accurately to 0.1 mg. The each artificial diet cube was cut 5cm on all sides; each pre weighed was placed in a petri dish. Helicoverpa armiger species exhibit high rates of cannibalism hence each petri dish contain individual larvae. Per weighted 4th instar larvae were allowed to feed on the test samples (i.e. plant extract and silver nanoparticles) containing 100 µg concentrations, while a control was maintained on artificial diet. The experiment was carried out in triplets. After 24 hours, the weights of diet cubes were recorded to track the larvae diet consumption. Mortality and morphological abnormalities linked with growth disrupting effects were identified in the larvae.

3.4. Extraction of midgut

After 24 hours of treatment, the midgut of 4th instar larvae was dissected under a microscope. The midgut was excised and rinsed in cold 50mM Tris-HCl. pH -7.4, then pulverized in a pestle and mortar in pH 7.4 Tris-HCl buffer at 4°C. The samples were centrifuged at 10,000g for 10 minutes in a 1.5ml eppendorf tube containing Tris-HCl buffer. The supernatant was precipitated with ammonium sulphate to achieve 70 per cent protein saturation. The precipitate was recovered by centrifugation at 10,000g for 10 minutes, diluted in 50 mM Tris-HCl buffer (pH 7.4), and kept at -20°C until used in further investigations.

3.5. RNA isolation

After the excavation of worms, they were stored in the nuclease free water and frozen in the -4^{0} C

until further process. On the day of RNA extraction the stored samples were washed mildly with the PBS to remove excess preserving solution and 200µl TRIZOl reagent (Invitrogen) was added for RNA extraction. Total RNA was isolated by homogenizing the samples using motorized pestle performing sonication for five strokes with 20secs "ON" and 15secs "OFF". The solution was combined with an equivalent volume of chloroform (200 µl) and centrifuged at 10,000 rpm for 10 minutes to separate it. The aqueous layer was collected and precipitated with isopropanol, and the particulate was reconstituted in nucleasefree water and quantified with Nanodrop after centrifugation (Denovix). Electrophoresis on a 1 per cent denatured agarose gel was used to assess the purity of the RNA. The quality of rRNA (28S and 18S) bands at a 2:1 ratio was used to assess RNA integrity.

3.6. cDNA Synthesis and real time PCR

Isolated RNA was used for cDNA synthesis using High capacity kit (Invitrogen). Briefly, the total reaction mixture was made up of buffer, dNTP's, random primers, reverse transcriptase enzyme with 1 μ g of RNA and total reaction volume was twenty microliters. The programme for cDNA synthesis was followed according the manufacturers manual. Upon completion of the synthesis the volume was made up to 100 μ l for the further analysis.

The cDNA were amplified for trypsin and Chymotrypsin (were coded as target 1 and 2 respectively) by real time PCR using Qiagen detection system with SYBR green as a detection dye. The specific primers for the target 1, 2 and beta actin were as follows,

Target1forward5'GCGTAAAGGATGCGGTTGG 3'

Reverse 5' CAGGATGGCAACCATCCATG 3'

Target2forward5'CACCATCTTCATCTTCCAATCCGTGTGC 3'

Reverse 5' GTGTTGATACGAGTACCACCGAAGAA C 3'

β-actin	forward	5'		
GATCGTGCGCGACATCAAG 3'				

Reverse 5' GCCATCTCCTGCTCGAAGTC 3'.

The SYBR green master mix containing cDNA products was amplified using the following protocol: denaturation at 95°C for 5 minutes, annealing at 60°C for 2 minutes, and extension at 72°C for 2 minutes. These conditions were repeated for 35 cycles and melting curve was obtained by heating the samples from 60°C to 95°C in 1°C increment in every cycle. The expression of target genes were normalized to the internal control β -actin and levels were calculated using delta Ct (2^{- $\Delta\Delta$ Ct}) method. The values were expressed as the relative fold gene expression. Here the trypsin and Chymotrypsin are named target 1 and target 2 respectively.

4. Results & Discussion

4.1 Characterization

XRD analysis: X-ray diffraction was used to determine the crystalline nature of biosynthesized AgNPs and the results are shown in figure 1(a-c). The peaks at 38.31° , 44.17° , 64.10° , and 77.56° in the LA sample correspond to planes (111), (200), (220), (311) of the cubic structure of silver nanoparticles, as shown in Figure 1(a). Particles in the nanoscale range are supported by the width and intensity of peaks. The (111), (200), (220), (311) planes of cubic phase of silver nanoparticles, respectively correspond to the 38.29°, 44.50°, 64.61°, 77.43° diffraction peaks for the CR sample (Fig.1b). The maxima of the CD sample at 38.43° , 44.64° , $64.810, 77.61^{\circ}$ correspond to the (111), (200), (220), and (311) planes of face centered cubic AgNp's, as shown in graph 1(c). The diffraction angles of all samples matches well with JCPDS data: 04-0783 [Kanniah et al., 2020; Adur et al., 2018]. The Scherrer equation, $D=K\lambda/\beta Cos\theta$, was used to determine the particle size of the samples. D is the crystallite size, and λ is the Xray beam wavelength β is the FWHM value. The calculated value of average crystallite size of LA, CR and CD sample is found to be 2.49nm, 2.37nm and 2.25 nm, respectively. As compared to LA and CD samples, the XRD spectrum of CR samples reveals a high crystallinity and a modest change in diffraction angle due to higher rate of reduction of AgNO₃.

EDX: The EDX spectra of samples LA, CR, and CD are shown in figure 2 (a-c). All samples have a prominent peak around 3 keV in their EDX spectra, indicating the existence of silver and justifying the formation of silver

nanoparticles. The weight percentage and atomic percentage of silver varies with different plant extracts due to the removal of unreacted organic and inorganic contaminants during product washing.

FTIR Analysis: FTIR spectroscopy was used to the different functional identify groups contained in samples and the results are displayed in figure 3 (a-c). The availability of flavonoids and terpenoids in the all plant allow for efficient silver extracts may nanoparticles capping and stability, resulting in the formation of all of these peaks. The peaks at 1611 cm⁻¹, 1271 cm⁻¹, 1065 cm⁻¹ are attributed to various squeezing vibrations of the OH, CH, C=C, and C-N groups, respectively [Kong et al., 2007]. The highest point 3078 cm⁻¹, 3327 cm⁻¹ due to chemisorbed/physisorbed H₂O and CO₂ molecules [Darezereshki, 2010]. The merging of peaks related to aldehydes (1692 cm^{-1}) , diketones (1566 cm⁻¹), alkanes (1469 cm⁻¹), amines (1334 cm⁻¹), and ethers (1185 and 1051 cm⁻¹). Peaks at 1593 cm⁻¹, 1509 cm⁻¹, and 1152 cm⁻¹ are attributed to the stretching of diketones, polysaccharides, and aromatic compounds, as seen in graph 3 (c). Peaks emerged in the mid all FTIR region of the spectrum at 2355.96 and atmospheric CO₂ 1358.68 cm⁻¹ due to adsorption [Sathyamoorthy and Mageshwari, 2012]. Weak absorption peaks associated with aromatic compounds (824.56 and 749.83 cm^{-1}), along with halogen compounds (538.93 cm⁻¹), were observed. The stretching vibration of aromatic chemicals included in extract causes a shift of strong peak at 825cm⁻¹ of CD sample towards lower wavenumber side as compared to spectra of LA and CR samples.

Raman spectroscopy: To probe the molecular structure and composition of samples we used the Raman spectroscopy and results are presented in figure 4 (a-c). Raman spectra of all samples show the slight change in pattern due to presence of aromatic and halogen compounds present in samples and their strong scattering properties. The sharp peaks around at 1348cm⁻¹ and 1550 cm⁻¹ due to the symmetric and antisymmetric vibration of C=O of carboxylic group [Chattopadhyay et al., 2005; Huang et al., 2008]. The small peaks noticed around at 1253cm⁻¹, 1145 cm⁻¹ 1064 and 932 cm⁻¹ are ascribed due to the in/out plane bending of C-C, C-N and C-H groups respectively [Chowdhury and Ghosh, 2007]. The peak at 1456 cm^{-1} is characteristic of asymmetric vibration of C-N group. Thus the change in these Raman bands, it

can be conclude that aromatic and halogen present in plant extract are encapsulated on surface of silver nanoparticles. The broad peak around at 2500cm⁻¹ for CD sample (Fig.4 (c)) due to vibration of aromatic and alkanes of extract absorbed on surface of plants nanoparticles. These results are well matches biosynthesis of AgNPs with by other researchers [Ajitha et al., 2014].

4.2. Role of AgNPs on *H.armigera* growth and development

Pesticide infiltration, often due to inadequate application, had long been a concern in the control of a variety of pests and weeds [Srinivasan et al.,2016] which seems to be particularly true for *H. armigera*, which is resistant to the majority of pesticides [Zaka et al.,2014]. Nanoparticles insecticidal properties are connected to physical qualities that drive physiological changes [Nel et al.,2006].

Under laboratory circumstances, the larvicidal effectiveness of LA, CR and CD crude extract and LA, CR, CD-Ag NPs was evaluated. After 24 hours of exposure with respective treatments, Helicoverpa armigera 4th instar larvae were assessed. All three plant extracts showed moderate Antifeedant activity and larval mortality, whereas no mortality was recorded for non-treated control. Silver nanoparticles had the highest mortality rate against larvae, while plant extracts had a moderate mortality rate, as shown in Fig.5. The above signifies that as compared to plant extract and control, larvae treated with AgNPs ingested the least amount of treated feed and died at a higher rate. The antifeedant activity data in Fig.6 demonstrated that the larvae feed on plant extract and silver nanoparticles reduced feeding activity compared control. Fig. to the 7(a-c) shows the physiological changes undergone by the larvae treated with silver nanoparticles i.e. decreased in larval body weight decreased body length and reduced movement. Fig7 (d-f) shows the physiological changes undergone by the larvae treated with crude extract i.e. body weight increased. Larvae fed AgNPs synthesised showed a substantial reduction in growth and development rate when compared to larvae fed plain extract and an experimental diet. Lepidoptera larvae have been observed to grow slowly when fed artificial more diets supplemented with additional metals such as zinc, cadmium, copper, and lead [Kramarz and Kafel, 2003; Stone et al., 2001]. The methanol

has been proven to be an excellent solvent for extracting bioactive compounds from plants with good larvicidal activity [Bagavan et al., 2009]. The detrimental impact of the plant extract might be likely to result in significant metabolites found in this plant [Isman, 2006; Shekari et al .,2008]. Synthesized AgNPs from E. prostrata have been shown to be efficient against the insect pest S. oryzae [Zahir et al., 2012]. AgNPs produced from Nelumbo nucifera leaves extract demonstrate larvicidal action versus Culex quinquefasciatus and Anopheles subpictus [Santhoshkumar et al., 2011]. Within 14 days of treatment, at a dosage of 20 ppm, the insecticidal effect of nanosilver colloid and ethanol-based colloid sulphur. Nano silver revealed nearly 100% mortality against casemaking clothes moth, T. pellionella (L.) larvae, damaging wool fibres [Ki et al., 2007]. The leaf dip method can be used to investigate larvicidal activity. Generally, the host plant's leaf discs are dipped different concentration in of manufactured nano biopesticides based on plant leaf extracts. After 24 hours of incubation with a consistent fresh supply leaves for good growth, the activity was assessed. After 24-96 hours of treatment, the mortality rates of the larvae are reported. CdS, nano-Ag, and nano-TiO2 nanoparticles have all been successfully tested against larvae [Chakravarthy et al., 2012].

4.3. Midgut protease (trypsin and Chymotrypsin) activity

Furthermore, our findings on gut proteases (trypsin and Chymotrypsin) support the produced silver nanoparticles from LA, CR and CD has an effective entomotoxicology potential on Helicoverpa armigera larvae. The LA and CR crude extract showed threefold increase in the activity while there was twofold decrease in respective synthesised silver nanoparticles. The CD extract showed fourfold increased and decrease in two fold in CD AgNPs. The consumption and survival rate of larvae in diets containing silver nanoparticles were significantly reduced. The AgNPs adhering to the target binding site resulted in a reduction in protease activity (trypsin and Chymotrypsin) and also suggest a link between both the protein and the nanoparticles. Bioinspired membranes can provide fundamental insights into the electrostatic interaction between nanoparticles and bio membranes by acting as excellent analogues to cell membranes [Lesniak et al., 2012]. Insect digestive proteases accelerate the release of free amino acids from ingested

Ishwarya M S, Lokesh S

protein, providing vital nutrients for growth and development. Many studies have been reported by researchers the activity of phytochemicals on the regions that control gastrointestinal motility and metabolism was identified as the cause of feeding restraint [Schmutterer, 1990]. The results of the quantitative real-time PCR studies revealed that all the crude plant extracts and sliver nanoparticles stimulated the protease activity. From fig. 8 and 9, indicated that the gut protease (trypsin and Chymotrypsin) activity slightly increased by four fold in the larvae treated with cured extract when compared to The larvae treated with silver control. nanoparticles showed three folds decreased activity when compared to crude extract. It was observed that amylase activity in *H. armigera* is modulated by biosynthesized silver nanoparticles [Kantrao et al., 2014]. The impact of rapid starch hydrolysis destruction was explored by studying the interaction of AgNPs with α -amylase [Ernest et al., 2012]. As pepsin and trypsin are so similar, it's logical to assume that AuNPs can improve trypsin stability as well. The pepsin anchored on AuNPs surface was more stable than just the free enzyme [Gole et al., 2001].

5.Conclusion:

We focused on environmentally friendly silver nanoparticle fabrication from LA, CR, and CD plant extracts in this study. Various approaches were used to characterise the physical properties of produced silver nanoparticles i.e. XRD spectra confirm the formation of AgNPs in phase centered cubic, FTIR results verify the formation and composition of components, and the findings of Raman spectroscopy are highly correlated. These silver nanoparticles were employed in control of pest Helicoverpa armigera. Despite crude plant extracts of LA, CR, and CD had no effect on larvae, nanoparticles synthesized from crude plant extract caused considerable cytotoxicity in the form of lower body weight and development in larvae. Hence silver nanoparticles synthesized from LA, CR and CD plant can effectively use as an agent to control Helicoverpa armigera pest (i.e. act as antifeedant agent, which subsequently deplete the growth of larvae and increase the mortality rate). Mimicking particles nature technique by enabling AgNPs as insecticidal chemicals could be a cost-effective way to manage insect pests while still minimizing the environmental impact. The above findings would also offer as a conceptual framework for the implementation of novel nanoparticles to combat pesticide resistance.

References:

- [1].De Oliveira JL, Campos EVR, Bakshi M, Abhilash PC, Fraceto LF. 2014. Application of nanotechnology for the encapsulation of botanical insecticides for sustainable agriculture: prospects and promises. Biotechnol. Adv. 32: 1550–1561.
- [2].Lade BD, Gogle DP, Nandeshwar SB. 2017. Nano-biopesticide to constraint plant destructive pests. J. Nanomed. Res. 6: 1–9.
- [3].Randall C, Hock W, Crow E, HudakWise C, Kasai J. 2013. National Pesticide Applicator Certification Core Manual. National Association of State Departments of Agriculture Research Foundation, Washington, DC, pp. 1–17.
- [4].Kamaraj C, Rahuman AA, Bagavan A. 2008. Screening for antifeedant and larvicidal activity of plant extracts against *Helicoverpa armigera* (Hübner), *Sylepta derogate* (F.) and *Anopheles stephensi* (Liston).Parasitol. Res. 103: 1361–1368.
- [5].Kamaraj C, Rahuman AA, Bagavan A.
 2008. Antifeedant and larvicidal effects of plant extracts against *Spodoptera litura* (F.), *Aedes aegypti* L. and *Culex quinquefasciatus* Say. Parasitol. Res. 103: 325–331.
- [6].Senthil-Nathan S. 2015. A review of biopesticides and their mode of action against insect pest's. In: Environmental Sustainability- Role of Green Technologies. Springer- Verlag, 49-63.
- [7].Sasson Y, Levy-Ruso G, Toledano O, Ishaaya I. 2007. Nanosuspensions: emerging novel agrochemical formulations. In: Ishaaya I, Nauen R, Horowitz AR (eds) Insecticides design using advanced technologies. Springer, Berlin. 1–32
- [8].Campolo O, et al., 2017. Citrus peel essential oil nanoformulations to control the tomato borer, Tuta absoluta: chemical properties and biological activity. Scientific Reports, 7 (1): 13036.
- [9].Gruère G, Clare N, Linda A. 2011. Agricultural, Food and Water Nanotechnologies for the Poor Opportunities, Constraints and Role of the Consultative Group on International Agricultural Research. J. Int. Food Policy Res. Inst. 1: 35.

- [10]. Worrall E, Hamid A, Mody K, Mitter N, Pappu H. 2018. Nanotechnology for plant disease management. Agronomy. 8(12):285
- [11]. Huang J, Li Q, Sun D, Lu Y, Su Y, Yang X, Wang H, Wang Y, Shao W, He N, Hong J and Chen C. 2007. Biosynthesis of silver and gold nanoparticles by novel sun dried *Cinnamomum camphora* leaf. Nanotechnology, 18:105104.
- [12]. Fouad H, Hongjie L, Hosni D, Wei J, Abbas G, Ga'al H, Jianchu M. 2018. Controlling *Aedes albopictus* and *Culex pipiens* pallens using silver nanoparticles synthesized from aqueous extract of *Cassia fistula* fruit pulp and its mode of action. Artif Cells Nanomed Biotechnol. 46:558–567.
- [13]. Srinivasan A, Giri AP and Gupta
 VS. 2006. Structural and functional diversities in lepidopteran serine proteases. Cell Mol Biol Lett. 11: 132–154.
- [14]. Sharma HC, Sharma KK, Seetharama N and Ortiz R. 2000. Prospects for using transgenic resistance to insects in crop improvement. Electron. J. Biotechnol. 3: 76–95.
- Swathi M, Mishra PK, Lokya V, [15]. Swaroop V, Mallikarjuna N, Dutta Gupta A. 2016. Purification and partial characterization of trypsin specific proteinase inhibitors from pigeonpea wild relative Cajanus platycarpus L. (Fabaceae) active against gut proteases of lepidopteran pest Helicoverpa armigera. Front. Physiol. 7:388.
- [16]. Patil CD, Borase HP, Suryawanshi RK, Patil SV. 2016. Trypsin inactivation by latex fabricated gold nanoparticles: a new strategy towards insect control. Enzym Microb Technol. 92:18–25.
- [17]. Santhoshkumar T, Rahuman AA, Bagavan A, Marimuthu S, Jayaseelan C, Kirthi AV, Kamaraj C, Rajakumar G, Zahir AA, Elango G, Velayutham K, Iyappan M, Siva C, Karthik L, Bhaskara Rao KV. 2012. Evaluation of stem aqueous extract and synthesized silver nanoparticles using *Cissus quadrangularis* against *Hippobosca maculata* and *Rhipicephalus* (Boophilus) *microplus*. Exp Parasitol. 132:156–165

Ishwarya M S, Lokesh S

- [18]. Rajakumar G, Rahuman AA. 2011. Larvicidal activity of synthesized silver nanoparticles using Eclipta prostrata leaf extract against filariasis and malaria vectors. Acta Trop. 118: 196–203.
- [19]. Kanniah P, Radhamani J, Chelliah P, Thangapandi Muthusamy N, EJJSB. Balakrishnan Thangapandi S, JR. Shanmugam R. 2020. Green synthesis of multifaceted silver nanoparticles using the flower extract of aerva lanata andevaluation of its biological and environmental applications. Chem.Select. 5: 2322-2331.
- [20]. Adur AJ, Nandini N, Shilpashree MK, Ramya R, Srinatha N. 2018. Biosynthesis and antimicrobial activity of silver nanoparticles using anaerobically digested parthenium slurry. J.Photochem. Photobio.B: Bio. 183: 30-34.
- [21]. Kong J,Yu S. 2007. Fourier transform infrared spectroscopic analysis of protein secondary structures. Acta biochimica et biophysica Sinica. 39: 549-559.
- [22]. Darezereshki E. 2010. Synthesis of maghemite (γ-Fe2O3) nanoparticles by wet chemical method at room temperature. Mater. Lett. 64:1471-1472.
- [23]. Sathyamoorthy R, Mageshwari K. 2012. Synthesis of hierarchical CuO microspheres: Photocatalytic and antibacterial activities. Physica E. 47:157-161.
- [24]. Chattopadhyay S, Lo HC, Hsu CH, Chen LC, Chen KH. 2005. Surfaceenhanced Raman spectroscopy using selfassembled silver nanoparticles on silicon nanotips. Chem.Mater. 17: 553-559.
- [25]. HuangY F, Chang HT, Tan W. 2008. Cancer cell targeting using multiple aptamers conjugated on nanorods. Anal.Chem. 80: 567-572.
- [26]. Chowdhury J, Ghosh M. 2007. Concentration-dependent surface-enhanced Raman scattering of 2-benzoylpyridine adsorbed on colloidal silver particles. J.Colloid. Inter.Sci. 277:121-127.
- [27]. Ajitha B, Ashok Kumar Reddy Y, Sreedhara Reddy P. 2014. Biosynthesis of silver nanoparticles using Plectranthus amboinicus leaf extract and its antimicrobial activity Spectrochimica Acta Part A. 128 : 257-262.

- [28]. VP, Srinivasan Nathan SS, Thanigaivel A, Edwin ES, Ponsankar A, Selin-Rani S, Pradeepa V. 2016. Developmental response of Spodoptera litura Fab. To treatments of crude volatile oil from Piper betle L. and evaluation of toxicity to earthworm, Eudrilus eugeniae Kinb. Chemosphere. 155:336-347.
- [29]. Zaka SM, Abbas N, Shad SA, Shah RM. 2014. Effect of emamectin benzoate on life history traits and relative fitness of Spodoptera litura (Lepidoptera: Noctuidae). Phytoparasitica. 42: 493-501.
- [30]. Nel A, Xia T, Madler L, Li N. 2006. Toxic potential of materials at the nanolevel. Science. 311, 622–627.
- [31]. Kramarz P, Kafel A. 2003. The respiration rate of the beet armyworm pupae (*Spodoptera exigua*) after multigeneration in toxication with cadmium and zinc. Environ. Pollut. 126, 1–3.
- [32]. Stone D, Jepson P, Laskowski R. 2011. Trends in detoxification enzymes and heavy metal accumulation in ground beetles (Coleoptera: Carabidae) inhabiting a gradient of pollution. Comp. Biochem. Physiol. Part C. 132, 105–112.
- [33]. Bagavan A, Kamaraj C, Elango G, Zahir AA, Rahuman AA. 2009. Adulticidal and larvicidal efficacy of some medicinal plant extracts against tick, fluke and mosquitoes. Vet Parasitol. 166(3–4):286– 292.
- [34]. Isman MB. 2006. Botanical insecticides, deterrents and repellents in modern agriculture and an increasingly regulated world. Annu Rev Entomol. 51:46– 66
- [35]. ShekariM, Sendi JJ, Etebari K, Zibaee A, Shadparvar A. 2008. Effects of Artemisia annua L. (Asteracea) on nutritional physiology and enzyme activities of elm leaf beetle. Xanthogaleruca luteola Mull. (Coleoptera: Chrysomelidae). Pest Biochem Physiol. 91:66–74.
- [36]. Zahir AA, Bagavan A, Kamaraj C, Elango G, Rahuman AA. 2012. Efficacy of plant-mediated synthesized silver nanoparticles against Sitophilus oryzae. J Biopest. 5:95–102

- [37]. Santhoshkumar T, Rahuman AA, Rajakumar G, Marimuthu S, Bagavan A, Jayaseelan, C. 2011. Synthesis of silver nanoparticles using Nelumbo nucifera leaf extract and its larvicidal activity against malaria and filariasis vectors. Parasitol. Res. 108: 693–702.
- [38]. Ki HY, Kim JH, Kwon SC, Jeong SH. 2007. A study on multifunctional wool textiles treated with nano-sized silver. J Mater Sci. 42:8020–8024.
- [39]. Chakravarthy AK, Chandrashekharaiah M, Khandakoor SB, Bhattacharyya A, Dhanabala K, Gurunatha K, Ramesh P. 2012. Bio efficacy of inorganic nanoparticles CdS, Nano-Ag and Nano-TiO₂ against Spodoptera litura (Fabricius) (Lepidoptera: Noctuidae). Curr. Biot. 6, 271–281.
- [40]. Lesniak A, Fenaroli F, Monopoli MP, Aberg C, Dawson KA, and Salvati A. 2012. "Effects of the presence or absence of a protein corona on silica nanoparticle uptake and impact on cells," ACS Nano, vol. 6, pp. 5845–5857.

- [41]. Schmutterer H. 1990. Properties and potential of natural pesticides from neem tree, *Azadirachta indica*, Annuval Review of Entomology. 35: 271-297.
- [42]. Kantrao S, Ravindra MA, Kamala Jayanthi PD, Venkataraman A. Modulatory effect of citrate reduced gold and biosynthesized silver nanoparticles on α -amylase activity. J. NanoparticlesID 2014; (829718):
- [43]. Ernest V, Shiny PJ, Mukherjee A, Chandrasekaran N. Silver nanoparticles: a potential nanocatalyst for the rapid degradation of starch hydrolysis by αamylase. Carbohydr, 2012; (352): 60–64.
- [44]. Gole A, Dash C, Ramakrishnan V, Sainkar SR, Mandale AB, Rao M, Sastry M. Pepsin-gold colloid conjugates: Preparation, characterization, and enzymatic activity. Langmuir 2001; 17 (5): 1674–1679.



Figure1. XRD spectra of Ag nanoparticles Graph (a-c) respectively, represent the XRD spectrum for LA, CR and CD samples.



Figure2. EDX spectra of AgNPs (a-c) respectively.



Figure3. FTIR spectra of sliver nanoparticles. Graph (a-c) represent FTIR spectrum of LA, CR and CD samples.



Figure4. Raman spectrometric analysis of AgNP's, Graph (a-c) represents the Raman spectra of LA, CR and CD samples.

International Journal of Agricultural Science http://iaras.org/iaras/journals/ijas





Fig.5. Effects of a diet containing LA, CR, and CD leaf extracts and LA, CR, CD AgNPs on the mortality rate of H. armigera larvae.



Fig.6. Effects Larvae body weight fed on diet containing LA, CR, CD crude leaf extracts and LA, CR, CD AgNPs.

Ishwarya M S, Lokesh S

International Journal of Agricultural Science http://iaras.org/iaras/journals/ijas



Fig.7. Influence of silver nanoparticles on larval weight (a-c) and crude leaf extract on larval weight (d-f).

	NOR	LA	CD	CR	LA AgNPs	CD AgNPs	CR AgNPs
TARGET 1 Ct	21.12	19.58	19.8	19.26	19.57	19.75	19.65
TARGET 2 Ct	16.21	15.01	14.7	14.8	15.01	15.04	15.01
Beta-actin Ct	15.12	15.42	15.86	15.06	14.09	15.04	14.04
T1 dcT (gene-ctrl)	6	4.16	3.94	4.2	5.48	4.71	5.61
T2 dcT (gene-ctrl)	1.09	-0.41	-1.16	-0.26	0.92	0	0.97
T1 2 ⁻ dct without	1	3.5801	4.169863	3.5801	1.433955248	2.445280555	1.31039340
T2 2 ⁻ dct without	1	2.828427	4.756828	2.549121	1.125058	2.12874	1.086735

Table1: Relative expression of traget1 (trypsin) and target 2 (Chymotrypsin) against Beta actin of leaf extracts and silver nanoparticles.



Fig.8. Expression of target 1 (trypsin) in the LA, CR, CD crude leaf extract and LA AgNPs, CR AgNPs, CD AgNPs.



Fig.9. Expression of target 2 (Chymotrypsin) in the LA, CR, CD crude leaf extract and LA AgNPs, CR AgNPs, CD AgNPs.

Abbreviations:

Helicoverpa armigera- H. armigera Fourier Transform infrared spectroscopy- FTIR X-ray diffraction- XRD Energy Dispersive X-Ray Spectroscopy-EDX *Leucas aspera* – *L. aspera*- LA

Cyperus rotundus- C. rotundus- CR

Cynodon dactylon- C. dactylon- CD

AgNPs- silver nanoparticles

Leucas aspera silver nanoparticles -LA-AgNPs

Cyperus rotundus silver nanoparticles - CR-AgNPs

Cynodon dactylon silver nanoparticles - CD-AgNPs

NBAIR- National Bureau of Agriculture Insect Resources.

Trypsin- Target 1

Chymotrypsin- Target 2