

























on yield contributing traits of Mash bean (*Vigna mungo* L.). *International Journal of Agronomy and Agricultural Research* **5**(6): 42-48.

*Food Science & Technology*, **4**:909-914.

[2]. Amrawat T, Solanki NS, Sharma SK, Jajoria DK and Dotaniya ML 2013. Phenology growth and yield of wheat in relation to agro meteorological indices under different sowing dates. *African Journal of Agricultural Research* **8**:6366-6374.

[3]. Biswas DK, Haque MM, Hamid A and Rahman MA 2002. Influence of plant population density on growth and yield of two Black Gram varieties. *Journal of Agronomy* **1**:83-85.

[4]. Jadhav PB, Kamble DR, Jadhav KT and Gadpale DL 2014. Performance of blackgram (*Vignamungo* (L.) Hepper) varieties to different sowing dates. *Advanced Research Journal of Crop Improvement* **5**(2):166-171.

[5]. Jondhale AN, Alse UN, Nirwal AD and Ghanwat PS 2018. Study of Agrometeorological Indices on Black Gram Cultivar under Varied Weather Condition. *International Journal of Current Microbiology and Applied Sciences* **7**(12): 2913-2919.

[6]. Karthick V and Mani K 2013. Perceptions on climate change and adaptations of dry land farmers in Tamil Nadu. *Environment and Ecology* **31**(3A):1540-1544.

[7]. Pal RK, Rao MNN and Murty NS 2013. Agro-meteorological indices to predict plant stages and yield of wheat for Foot Hills of Western Himalayas. *International Journal of Agriculture*