

- [6] C. Beale, "The Behavioral Ecology of Disturbance Responses", Escholarship.org, 2019. [Online]. Available: <https://escholarship.org/uc/item/43m7b2d5>. [Accessed: 09- May- 2019].
- [7] M. Yoklavich, M. Love and K. Forney, "A fishery-independent assessment of an overfished rockfish stock, cowcod (*Sebastes levis*), using direct observations from an occupied submersible", Canadian Journal of Fisheries and Aquatic Sciences, vol. 64, no. 12, pp. 1795-1804, 2007. Available: 10.1139/f07-145 [Accessed 9 May 2019].
- [8] Fao.org, 2019. [Online]. Available: <http://www.fao.org/3/a-a0210e.pdf>. [Accessed: 09- May- 2019].
- [9] F. Uiblein, P. Lorance and D. Latrouite, "Behaviour and habitat utilization of seven demersal fish species on the Bay of Biscay continental slope, NE Atlantic", Marine Ecology Progress Series, vol. 257, pp. 223-232, 2003. Available: 10.3354/meps257223 [Accessed 9 May 2019].
- [10] V. TRENKEL, P. LORANCE and S. MAHEVAS, "Do visual transects provide true population density estimates for deepwater fish?", ICES Journal of Marine Science, vol. 61, no. 7, pp. 1050-1056, 2004. Available: 10.1016/j.icesjms.2004.06.002 [Accessed 9 May 2019].
- [11] W. Beebe, "Preliminary Account of Deep Sea Dives in the Bathysphere with Especial Reference to One of 2200 Feet", Proceedings of the National Academy of Sciences, vol. 19, no. 1, pp. 178-188, 1933. Available: 10.1073/pnas.19.1.178 [Accessed 9 May 2019].
- [12] C. Wallington, "The Face of the Deep. B. C. Heezen and C. D. Hollister, Oxford University Press, New York, 1971. 220 x 2X5 mm., xii and 659 pages (with indexes), 3 tables, 220 maps and diagrams, 355 photographic plates (8 in colour). Cloth \$29.95, paper \$12.50.", Cartography, vol. 9, no. 1, pp. 48-49, 1975. Available: 10.1080/00690805.1975.10437848 [Accessed 9 May 2019].
- [13] Churashima.okinawa, 2019. [Online]. Available: <https://churashima.okinawa/userfiles/files/page/ocrc/p5-21.pdf>. [Accessed: 09- May- 2019].
- [14] P. Lorance and V. Trenkel, "Variability in natural behaviour, and observed reactions to an ROV, by mid-slope fish species", Journal of Experimental Marine Biology and Ecology, vol. 332, no. 1, pp. 106-119, 2006. Available: 10.1016/j.jembe.2005.11.007 [Accessed 9 May 2019].
- [15] C. Szegedy et al., "Going Deeper with Convolutions", arXiv.org, 2019. [Online]. Available: <https://arxiv.org/abs/1409.4842>. [Accessed: 08- Mar- 2019].
- [16] A. Hollowed et al., "Projected impacts of climate change on marine fish and fisheries", ICES Journal of Marine Science, vol. 70, no. 5, pp. 1023-1037, 2013. Available: 10.1093/icesjms/fst081 [Accessed 11 May 2019].
- [17] M. Turk and A. Pentland, "Eigenfaces for Recognition", Journal of Cognitive Neuroscience, vol. 3, no. 1, pp. 71-86, 1991. Available: 10.1162/jocn.1991.3.1.71 [Accessed 8 May 2019].
- [18] L. Meng, T. Hirayama and S. Oyanagi, "Underwater-Drone With Panoramic Camera for Automatic Fish Recognition Based on Deep Learning", IEEE Access, vol. 6, pp. 17880-17886, 2018. Available: 10.1109/access.2018.2820326 [Accessed 10 March 2019].