

parallel state) is favored compared to the antiparallel, antiferromagnetic state

According to theory and available data [4,8]

$$J = J_0 \exp(-\beta r) \quad (2)$$

where J_0 is the exchange integral at Van der Waals distance, r is the spin-spin distance and β is the distance decay factor

In many systems, spin exchange takes place by superexchange through a bridge or a through space interaction mediated by the molecules of the intervening medium:



Scheme 1.2 Spin exchange via a bridge

Anderson [9] in his pioneering work used a perturbational approach to relate the singlet-triplet splitting of a two-spin system to the magnitude of the electron-transfer superexchange coupling between the radical ion pair (RP) state and surrounding states n and that state to which it is coupled at the nuclear coordinate of the relaxed RP state. Then McConnell [6] has shown that the electronic coupling matrix element (H_{AB}) between two redox centers separated by a covalent bridge composed of n identical repeat units depends on the coupling strength between the redox sites and the bridge (h_{Ab} , h_{bB}), the coupling between adjacent bridge elements (h_{bb}), and the tunneling energy gap ($\Delta\varepsilon$). The latter was defined as the virtual energy required to remove an electron from the donor, or a hole from the acceptor, and place it on the bridge.

According to theory presented in [10]

$$H_{DA} = \frac{h_{Db}}{\Delta\varepsilon} \left(\frac{h_{bb}}{\Delta\varepsilon} \right)^{n-1} h_{bA} \quad (3)$$

2.2 Electron Transfer

Electron transfer (ET) is one of the most ubiquitous and fundamental phenomena in

Nature. ET is found to be a key elementary step in many important processes in chemistry, biology and physics including systems of the light energy conversion.[1-7.]

Electron transfer (ET), as in any other chemical reaction, is accompanied by a change in the nuclear and electronic configurations and is described by the Fermi's golden rule equation for a transition with rate constant k_{tr} [11]

$$k_{tr} = \frac{J_{ex}^2}{h} FC(4)$$

which includes the exchange integral (an electronic coupling term) J_{ex} depending on the overlap of electronic wave functions in the initial and final state of the process; FC is the Franck-Condon factor related to the probability of vibrational and translational transition states accounting for the effects of nuclear motion. FC factor related to the much slower changes in the nuclear configuration, whereas the exchange integral depends on the instantaneous electronic configuration in the Born-Oppenheimer approximation.

According to the fundamental Marcus two state model [11] the distortion of the reactants, products and solvent from their equilibrium configuration is described by identical parabolas in reaction coordinate space for the initial and final states. Within the adiabatic regime (strong electronic coupling, where the resonance integral J_{ex} , $V > 200 \text{ cm}^{-1}$), the value of the electron transfer rate constant is given by the Marcus-Eyring theory of the transition state:

$$k_{ET} = \frac{k_B T}{h} \exp - \left[\frac{(\lambda + \Delta G_0)^2}{4\lambda k_B T} \right] \quad (5)$$

where λ is the reorganization energy defined as the energy for electron excitation without distortion of the nuclear frame and $-\Delta G_0$ is the process thermodynamic driving force

The theory developed by Levich, Dogonadze [13] was related to non-adiabatic electron transfer between donor (D) and acceptor (A) centers. The theory is based on Fermi's Golden Rule (Eq. 1), the Landau-Zener Eq. [14,15] and the Marcus formula (Eq. 5). Using these

concepts, the authors proposed a formula for non-adiabatic ET

$$k_{ET} = \frac{2\pi V^2}{h\sqrt{4\pi\lambda k_B T}} \exp\left[-\frac{(\lambda + \Delta G_0)^2}{4\lambda k_B T}\right] \quad (6)$$

A principle connection between exchange integral and electron transfer coupling factor was established by Anderson [9]. In the Anderson equation for a radical (ion-radical) pair (RP): the indicated matrix elements couple the singlet and triplet RP states to states n , E_{RP} and E_n are energies of these states, respectively, and λ is the total nuclear reorganization energy of the charge-transfer reaction:

$$2J = \Delta E_S - \Delta E_T = \left[\sum_n \frac{|\langle \Psi_{RP} | V_{RP-n} | \Psi_n \rangle|^2}{E_{RP} - E_n - \lambda} \right]_S - \left[\sum_n \frac{|\langle \Psi_{RP} | V_{RP-n} | \Psi_n \rangle|^2}{E_{RP} - E_n - \lambda} \right]_T \quad (7)$$

were Ψ is the corresponding wave function. The Anderson approach was effectively exploited [16-18].

3. Semi-empirical methods of calculation of ET rate constants

A semi-empirical approach for the quantitative estimation of the effect bridging the group on the long-distance electron transfer LDET was developed by Likhstenshtein [3,4,7,19].

The approach is based on analogy between superexchange in electron transfer (ET) and such electron exchange processes as triplet-triplet energy transfer (TTET) and spin-exchange (SE). The spin exchange and TT phenomena may be considered an idealized model of ET without or with only a slight replacement of the nuclear frame.

As was shown in [19], experimental data on the dependence of the rate constant of triplet triplet energy transfer k_{TT} and the exchange integral J_{SE} on the distance between the centers (ΔR) lies on two curves, which are approximated by the following equation:

$$k_{TT}, J_{SE} \propto \exp(-\beta \Delta R) \quad (8)$$

It was found that for systems in which the centers are separated by a “non-conductive” medium (molecules or groups with saturated chemical bond) $\beta_{TT} = 2.6 \text{ \AA}^{-1}$. For systems in which the radical centers are linked by “conducting” conjugated bonds, β_{SE} is 0.3 \AA^{-1} . The data on exchange integrals for paramagnetic center separated by “non conducting” media were obscured.

Above mentioned analogy can allow to predict corresponding distance dependent for SE and therefore for ET. We suggest that in the first approximation

$$V_{ET}^2 J_{SE} J_{TT} \propto \exp(-n\beta R) \quad (9)$$

where $(|V_{CR}|^2)$ is the square of the resonance integral, J_{SE} is the exchange integral, J_{TT} is the triplet-triplet transfer integral, R is the distance between the interacting centers and β_i is a coefficient which characterizes the degree of the integral decay (decay factor). In the first approximation, $n = 2$ for the ET and SE processes with the overlap of two orbitals and $n = 4$ for the TT process in which four orbitals overlap, that is, ground and triplet states of the donor and ground and triplet states of the acceptor).

The ratios

$$\gamma = \frac{J_{SE}^0}{J_{SE}^x} \quad (10)$$

were J_{SE}^0 and J_{SE}^x are the exchange integrals for two non-bridged radicals and radicals tethered by a media x , can be considered as attenuation factor of the exchange interaction of SE through the given bridge. Taking into account Eq. 8.3 with value $n = 2$ for SE and ET, an expression for the dependence of the attenuation parameters for SE and ET on the distance between remote donor and acceptor centers $D_R D_A$ can be given as Eq. 8.2 with $\beta_{ET}(\text{nc}) = 0.5\beta_{TT} = 1.3 \text{ \AA}^{-1}$ for a “non-conducting” medium and $\beta_{ET}(\text{c}) = 0.3 \text{ \AA}^{-1}$ for a “conducting” bridge. The value of $\square_{ET} =$

1.3 \AA^{-1} is found to be close to that obtained by analysis of k_{ET} , which is proportional to $|V_{CR}|^2$, on the distance ΔR in model and biological systems (Fig.1).

Similar but not identical approach was developed in [16]. The authors also paid attention on similarity of long-range intramolecular electron transfer and positive ions (hole) transfer, and triplet energy transfer and showed that $\beta^{TT}/\beta^{ET} = 2$.




Figure 1 The observed linear dependence of the logarithm of the rate constant of the electron transfer in RCs of purple bacteria and plant photosystem I ($\log k_{ET}$) on the edge-edge distance between the donor and acceptor centers (r_{DA}) (circles), solid line is the dependence predicted by Eqs. 9-10 for the exchange integral J_{SE} [19].

The empirical data on the exchange integral values (J_{ET}) for the spin-spin interactions in systems with known structure, that is, biradicals, transition metal complexes with paramagnetic ligands and monocystals of nitroxide radicals have served as a basis for the estimation of the value of the attenuation parameter γ_{\square} for the exchange interaction through a given group X [19]. By our definition, the γ_{\square} is

$$\gamma_X = \frac{J_{RYZP}}{J_{RYXZP}} \quad (11)$$

where R is a nitroxide or organic radical, P is a paramagnetic complex or radical and X, Y, and Z are chemical groups in the bridge between R

and P. Table 1 shows the results of the calculation of parameter γ_X from empirical data on the spin exchange in the nitroxidebiradicals by Eq. 11

Table 1. Attenuation parameter γ_X calculated from empirical data on the spin exchange in nitroxidebiradicals and complexes with paramagnetic ligands: γ_X for individual group, γ_{hb} for hydrogen bond and γ_v for Van der Waals contact.[19].

Group, X	γ_X	Group, X	γ_X
C ₆ H ₄	6.00±0.03		
C=C	1.7	-NH-CO-	55
C=O	8.4±0.4	γ_v	50
C			
NH	6.5	γ_{hb}	10
O	5	H	12
S=O	2.1	SO ₂	2.2
	3.5	RP=O	2.40±0.03

Thus, the experimental dependence of exchange parameters k_{TT} and J_{SE} on the distance between the exchangeable centers and the chemical nature of the bridge connecting the centers may be used for evaluating such dependences for the resonance integral in the ET equations (Eqs.9-10).

4. Estimation of spin exchange integral and ET rate constants in donor –acceptor pairs

Data presented in Table 1. and Eq. 9-10 was used for the analysis of alternative electron transfer pathways in chemical and biological systems. The following equations for the spin exchange integral and rate constant of electron transfer in donor-acceptor pairs are suggested:

$$J_{SE} = J_0 \rho_{S1} \rho_{S2} \prod_i \gamma_i^{-1} \dots \dots \dots \quad (12)$$

and

$$k_{ET} = k_0 \rho_D \rho_A \prod_i \gamma_i^{-1} \quad (13)$$

where $J_0 = 10^{14} \text{ s}^{-1}$ exchange integral at Van der Waals contact, $\rho_{S1,2}$ - spin density at Van der Waals contact, γ_i - attenuation factors, $k_0 = 5 \times 10^{10} \text{ s}^{-1}$

is the rate constant of ET in a nonbridged DA pair Eqs. 12 and 13 prediction of values of exchange integral and ET rate constant $J_X / k_X = J_0 / k_0 = 200$.

Values of J_{SE} and k_{ET} calculated by Eqs. 12 and 13 were compared with corresponding experimental data. For rate constant of recombination of charge separated primary and secondary quinons [17] (Fig. 2) in the photosynthetic reaction centers, employment Eq. 13 led to following calculated value: calculated (this work) $k_{ETqq} = .5 \times 10^8 \text{ s}^{-1}$ as compare with $k_{ETqq} = .9 \times 10^8 \text{ s}^{-1}$ calculated by Calvo *et al.* and experimental $k_{ETqq} = .3 \times 10^8 \text{ s}^{-1}$ [17].





Figure 2 Segment of electron transfer chain of bacterial photosynthetic center [17]

Eq. 12 with values $J_0 \approx 10^{14} \text{ s}^{-1}$, $\rho_{S1} = \rho_{S2} = 0.2$ and $\gamma_1 \gamma_2 = 36$ for the donor-acceptor pair [18]:



I

and two phenyl bridged groups gave calculated $J_2 = 10^{10} \text{ s}^{-1}$ as compared with experimental $J_2 = 6 \times 10^9 \text{ s}^{-1}$ [18] A good correlation between experimental J values ($n = 2-5$) and that predicted by Eq. 8.8 take place (Fig. 3)




Figure 3 Logarithmic plot of the exchange integral $2J$ vs donor-acceptor distance, r_{DA} . Points experimental data [27], solid line predicted from Eq. 12 [18]

For the rate constant of charge separation in with $n=1$, Eq. 8.9 predicted $k_{ICS} = 3 \times 10^{10} \text{ s}^{-1}$ versus experimental $k_{ICS} = 5 \times 10^{10} \text{ s}^{-1}$

As one can see from Fig. 4, for compound I with $n = 1-3$ experimental data of the charge recombination rate constant, k_{CS} vs donor-acceptor distance fit to the dependence predicted by Eq. 13, while the experimental dependence rate constant of charge separation is steeper than for predicted one. This discrepancy can be explained suggesting that the spin exchange and recombination involve the same triplet orbitals, while charge separation occurs from singlet orbital.






Figure 4 (A) Logarithmic plot of the charge separation rate constant k_{CS} vs donor-acceptor distance, r_{DA} . (B) Logarithmic plot of the charge recombination rate constant, k_{CR} vs donor-acceptor distance, r_{DA} , solid line predicted from Eq. 13. [18].


Probing protein environment with dual fluorophore-nitroxide (FNO \cdot) molecules in which fluorophore is tethered with nitroxide, a fluorescence quencher opens unique opportunities to study molecular dynamics and micropolarity of a protein medium, intramolecular fluorescence quenching (IFQ), electron transfer, photoreduction and light energy conversion [3,4,7,20-23].

The following scheme describes photophysical and photochemical processes in the dual molecule FN I:



(FN1)

incorporated into bovine serum albumin were investigated (Fig. 5)



Scheme 2 Light energy conversion in the dual fluorophore-nitroxide (FNO \cdot) molecules

The fluorophore and nitroxide segments of the probes allow monitoring of molecular dynamics

and also make it possible to measure micropolarity of the medium in the vicinity of the donor (by fluorescence technique) and acceptor (by ESR) moieties. Exchange, energetic and molecular dynamic factors affecting photoinduced electron transfer in a donor-acceptor pair (D-A) incorporated into bovine serum albumin were investigated (Fig. 5)[20-23] The kinetics of reversible and irreversible intramolecular electron transfer from the donor to the acceptor were monitored by picoseconds time resolved fluorescent and ESR techniques corresponding at temperature range from 70 K to 300 K.

For the fluorescent-nitroxide probe FN1, Eq.13 related to ET along the bridge chain, after taking $ak_0 = 5 \times 10^{12} \text{ s}^{-1}$, $\rho_1 = \rho_2 = 0.4$ and the corresponding attenuation factors γ_i picked up from Table 1, gave $k_{ET} = 1.1 \times 10^8 \text{ s}^{-1}$, while the experimental value was found as $k_{ET}(\text{exp}) = 1.0 \pm 0.1 \times 10^8 \text{ s}^{-1}$ [22].

For the fluorescent-nitroxide probe FN 1, Eq. for ET through the space, after taking $k_0 = 5 \times 10^{12} \text{ s}^{-1}$, $\rho_1 = \rho_2 = 0.4$ and $\beta = 1.3 \text{ \AA}^{-1}$, gives the value of $k_{ET} = 8.8 \times 10^7 \text{ s}^{-1}$. For ET along the bridge chain, using Eq.13 with $\rho_1 = \rho_2 = 0.4$ and attenuation factors for phenyl segment $\gamma_i = 0.4$ (Table 1) led to $k_{ET} = 1.1 \times 10^8 \text{ s}^{-1}$, while the experimental value was found as $k_{ET}(\text{exp}) = 1.0 \pm 0.1 \times 10^8 \text{ s}^{-1}$.




Figure 5 Dual fluorophore-nitroxide (FNO[•]) molecule incorporated in the binding center of albumin.

Conclusion

Available theoretical considerations and experimental data have unambiguously proved a connections between electron transfer, spin exchange and triplet triplet energy transfer. When in these processes the same orbitals are involved, this connection is quantitative one. The suggested semiempirical approach allowed to estimate values of exchange integral and electron transfer rate constant are in fairly good agreement with corresponding experimental data and can be used for an estimation these values in unknown system of interest.

References

[1] Bixon M. and Jortner J. Electron Transfer from Isolated Eolecules to Biomolecules, in Jortner, J and Bixson, M. (eds.), *Advances in Chemical Physics*, 107, Parts 1 and 2, John Wiley & Sons, NY, 1999, pp. 35-202.

[2] Marcus R. A., Electron Transfer Past and Future, in Jortner, J. and Bixson, M. (eds.), *Advances in Chemical Physics*, 107, Parts 1 and 2, John Wiley & Sons, NY, 1999, pp.13-19,

[3] Likhstenshtein G. I., *Solar Energy Conversion. Chemical Aspects*, WILEY-VCH, Weinhem, 2012.

[4] Likhstenshtein G. I. *Electron Spin in Chemistry and Biology: Fundamentals, Methods, Reactions Mechanisms, Magnetic Phenomena, Structure Investigation*. Springer 2016

[5] Likhstenshtein G. I., Syrtsova L. A., Samuilov V.D., Frolov E. N., Borisov A.U., Bogatyrenko V. R., Photosynthetic reaction center. *XII International Botanic Congress:*

Theses of Reports, (Nauka, Leningrad, 1975, p.429.

[6] Likhstenshtein G. I., Kotel'nikov A. I., Kulikov A. V., Syrtsova L. A., Bogatyrenko V. R., Mel'nikov A. I., Frolov E. N., Berg A. J., Some peculiarities of the electronic transfer in redox enzymes. *Intern J Quant Chem* 1979, 16, 419-435.

[7] Likhstenshtein G. I., *New Trends In Enzyme Catalysis and Mimicking Chemical Reactions*, N.Y. Kluwer Academic/ Plenum Publishers, 2003.

[8] Zamaraev, K. I, Molin, Y. N., Salikhov K. M. *Spin Exchange. Theory and Physicochemical application*, Springer-Verlag, Heidelberg, 1981.

[9] Anderson P. W., New Approach to the Theory of Superexchange Interactions" *Physical Review*, 1959, 115, 2-15.

[10] McConnell H. M., Intramolecular Charge Transfer in Aromatic Free Radicals, *Journal of Chemical Physics*, 1961, 35, 508-515.

[11] Marcus R. A. Transfer Reactions in Chemistry. Theory and Experiment, *Pure and Applied Chemistry*, 1997, 69, 13-29.

[12] Neil R., Kestner N. R., Logan J., Jortner J., Thermal Electron Transfer Reactions in Polar Solvents" *Journal of Physical Chemistry*, 1974, 78, 2148-2166.

[13] Levich V. G., Dogonadze, R., Quantum Mechanical Theory of Electron transfer in Polar Media, *Doklady Akademii Nauk*, 1959 78, 2148-2153.

[14] Landau, L. Zur Theorie der Energie Uebertragung, *Phys Zur Sovietunion*, 1932, 2, 46-51. [15] Zener C. Dissociation of Excited Molecules by External Perturbation'. *Proceeding Royal Society A*, 1933, 140:660- 668.

[16] Closs G. L, Johnson M. D, Miller J. R., Piotrowiak P., A Connection Between Intramolecular Long-range Electron, Hole, and Triplet Energy Transfers" *P. Journal of American Chemical Society* 1989. 111:3751-3753.

[17]. Calvo R., Abresch E. C., Bittl R., Feher G, Hofbauer W., Isaacson R. A. Lubitz W., Okamura M. Y., Paddock, M. L. EPR Study of the Molecular and Electronic Structure of the Semiquinone Biradical

$Q_A^-Q_B^-$ in Photosynthetic Reaction Centers from *Rhodobactersphaeroides*, *Journal of American Chemical Society*, 2000, 122, 7327-7341.

[18]. Weiss E.A., Ahrens M. J., Sinks L., Gusev E. A., Ratner M. A., Wasielewski, M. R., Making a Molecular Wire: Charge and Spin Transport through *para*-Phenylene Oligomers, *Journal of American Chemical Society* 2004 126, 5577-5586

[19]. Likhtenshtein G. I., Role of Orbital and Dynamic Factors in Electron Transfer in Reaction Centers of Photosynthetic Systems, *Journal of Photochemistry and Photobiology A: Chemistry* 1996, 96, 79-92.

[20] Likhtenshtein, G. I., Ishii, K., and Nakatsuji, "Dual Chromophore-Nitroxides: Novel Molecular Probes, Photochemical and Photophysical Models and Magnetic Materials", *Photochemistry and Photobiology*, 2007, 83, 871-881.

[21] Likhtenshtein G. I., Yamauchi J., Nakatsuji, S., Smirnov A Tamura, R., *Nitroxides: Application in Chemistry, Biomedicine, and Materials Science*. WILEY-VCH, Weinheim 2008.

[22] Likhtenshtein G. I. Factors Affecting Light Energy Conversion in Dual Fluorophore-Nitroxide Molecules in Solution and a Protein, *Pure and Applied Chemistry*, 2008, 80, 2125-2139.

[23] Likhtenshtein G. I., Pines, D., Pines, E., Khutorsky, V. Effect of a Protein on the Light Energy conversion in Dual Fluorophore-nitroxide Probes Studied by ESR and Fluorescence Spectroscopy, *Applied Magnetic Resonance* 2009, 35, 459-472.