















- [6] JH. Lee, KG. An, “Integrative restoration assessment of an urban stream using multiple modeling approaches with physical , chemical and biological integrity indicators”, Ecological Engineering, vol. 62, pp. 153-167, 2014.
- [7] AK. Stefanie, L. Brian, R. Anthony, “Using ecological niche models to predict the abundance and impact of invasive species: application to the common carp”, Ecological Applications, vol. 2, pp. 203-213, 2011.
- [8] JP. Steven, PA. Robert, EC. Robert “Maximum entropy modeling of species geographic distributions”, Ecological Modelling, vol. 190, pp. 231-259, 2006.
- [9] EE. Prepas, FA. Rigler, “Improvements in qualifying the phosphorus concentration in lake water”, Canadian Journal of Fisheries and Aquatic Sciences, vol. 39, pp. 822-829, 1982.
- [10] American Public Health Association, “Standard methods for the examination of water and wastewater”, 19th Ed. American Public Health Association. Washington, DC, 1991.
- [11] MT. Barbour, J. Gerritsen, JB. Stribling, “Rapid bioassessment protocols for use in streams and waderable rivers: periphyton, benthic macroinvertebrates, and fish”, Second Edition, EPA 841-B-99-002. US EPA; Office of Water, Washington, 1999.
- [12] K. Takamura, “Performance as a fish predator of largemouth bass (*Micropterus salmoides* -Lacepede) invading Japanese freshwaters; a review”, Ecological resources, vol. 22, pp. 940-946, 2007.
- [13] AN. Fedoruk, “Feeding relationship of Walleye and smallmouth bass”, Journal of the Fisheries Board of Canada, vol. 23, pp. 941-943, 1966.